

FILE 'REGISTRY' ENTERED AT 14:49:23 ON 22 SEP 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 21 SEP 2006 HIGHEST RN 908228-18-2
DICTIONARY FILE UPDATES: 21 SEP 2006 HIGHEST RN 908228-18-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 30, 2006

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>

Uploading C:\Program Files\Stnexp\Queries\150f.str

L7 STRUCTURE UPLOADED

=> d l7

L7 HAS NO ANSWERS

L7 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation.

=> s l7 sss sam

SAMPLE SEARCH INITIATED 14:49:48 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 27 TO ITERATE

100.0% PROCESSED 27 ITERATIONS

0 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 229 TO 851

PROJECTED ANSWERS: 0 TO 0

L8 0 SEA SSS SAM L7

=> s l7 sss full

FULL SEARCH INITIATED 14:50:03 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 755 TO ITERATE

100.0% PROCESSED 755 ITERATIONS

20 ANSWERS

SEARCH TIME: 00.00.01

L9 20 SEA SSS FUL L7

=> fil hcaplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	166.94	169.73

FILE 'HCAPLUS' ENTERED AT 14:50:09 ON 22 SEP 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 22 Sep 2006 VOL 145 ISS 14
FILE LAST UPDATED: 21 Sep 2006 (20060921/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 19

L10 19 L9

=> d 110 ibib hitstr abs all

L10 ANSWER 1 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2006:165092 HCAPLUS

DOCUMENT NUMBER: 144:370172

TITLE: New C2- and C1-Symmetric phosphorus ligands based on carbohydrate scaffolds and their use in the iridium-catalysed hydrogenation of ketimines

AUTHOR(S): Guiu, Ester; Aghmiz, Mohamed; Diaz, Yolanda; Claver, Carmen; Mesequer, Benjami; Militzer, Christian; Castillon, Sergio

CORPORATE SOURCE: Departament de Quimica Analitica i Quimica Organica, Universitat Rovira i Virgili, Tarragona, 43005, Spain

SOURCE: European Journal of Organic Chemistry (2006), (3), 627-633

CODEN: EJOCFK; ISSN: 1434-193X

PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 144:370172

IT 666825-71-4


RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)

RN 666825-71-4 HCAPLUS

CN D-Mannitol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]-
(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



IT 666826-33-1P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)

RN 666826-33-1 HCAPLUS

CN D-Mannitol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]-, 3-[bis(2,4-dimethylphenyl)phosphinite] (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



AB D-Mannitol-derived C2-sym. diarylphosphinite and C1-sym. diaryl phosphite-phosphinite ligands were prepared from silylated D-glucosamine; asym. hydrogenation of acetophenone benzylimine, catalyzed by iridium complexes with new ligands gave N-benzyl-1-phenylethylamine with 73% ee. The C2-sym. diphosphinites, (3R,4R)-2,5-(TBDPSO)2-3,4-(Ar₂PO)-tetrahydrofurans (10a-d; Ar = Ph, 4-MeOC₆H₄, 4-CF₃C₆H₄, 3,5-Me₂C₆H₃) were prepared by reaction of (3S,4S)-2,5-(TBDPSO)2-3,4-tetrahydrofurandiol (12) with Ar₂PCl or Ar₂PNEt₂; the mono-substituted (3R,4S)-2,5-(TBDPSO)2-4-(Ar₂PO)-3-tetrahydrofuranol was esterified by 2,2'-methylenebis[4-methyl-6-CMeR1R2-phenyl] phosphorochloridites to give the corresponding C1-sym. phosphite-phosphinites [11a,b, R₁ = R₂ = Me, R₁+R₂ = (CH₂)₅]. Various procedures for synthesizing the phosphinite function were explored in order to improve the yield of the reaction. Results were best when Ph₂PNEt₂ was used in the presence of tetrazol as catalyst. The prepared ligands, which have different electron-donating or electron-withdrawing aryl groups were added to iridium complexes producing catalyst precursors active in the asym. hydrogenation of acetophenone N-benzyl- and N-phenylimines (17, 19). Cationic iridium complexes were more active than the neutral analogs. The use of additives was, in general, detrimental to both the conversion and the enantioselectivity. In the hydrogenation of 17, results were best with ligand 11a (76% ee), but in the hydrogenation of 19 (70% ee) they were best with ligand 10b.

AN 2006:165092 HCAPLUS

DN 144:370172

ED Entered STN: 23 Feb 2006

TI New C2- and C1-Symmetric phosphorus ligands based on carbohydrate scaffolds and their use in the iridium-catalysed hydrogenation of ketimines

- AU Guiu, Ester; Aghmiz, Mohamed; Diaz, Yolanda; Claver, Carmen; Meseguer, Benjami; Militzer, Christian; Castillon, Sergio
- CS Departament de Química Analítica i Química Orgànica, Universitat Rovira i Virgili, Tarragona, 43005, Spain
- SO European Journal of Organic Chemistry (2006), (3), 627-633
CODEN: EJOCFK; ISSN: 1434-193X
- PB Wiley-VCH Verlag GmbH & Co. KGaA
- DT Journal
- LA English
- CC 29-7 (Organometallic and Organometalloidal Compounds)
Section cross-reference(s): 28, 33
- OS CASREACT 144:370172
- AB D-Mannitol-derived C2-sym. diarylphosphinite and C1-sym. diaryl phosphite-phosphinite ligands were prepared from silylated D-glucosamine; asym. hydrogenation of acetophenone benzylimine, catalyzed by iridium complexes with new ligands gave N-benzyl-1-phenylethylamine with 73% ee. The C2-sym. diphosphinites, (3R,4R)-2,5-(TBDPSO)2-3,4-(Ar2PO)-tetrahydrofurans (10a-d; Ar = Ph, 4-MeOC6H4, 4-CF3C6H4, 3,5-Me2C6H3) were prepared by reaction of (3S,4S)-2,5-(TBDPSO)2-3,4-tetrahydrofurandiol (12) with Ar2PCl or Ar2PNET2; the mono-substituted (3R,4S)-2,5-(TBDPSO)2-4-(Ar2PO)-3-tetrahydrofuranol was esterified by 2,2'-methylenebis[4-methyl-6-CMeR1R2-phenyl] phosphorochloridites to give the corresponding C1-sym. phosphite-phosphinites [11a,b, R1 = R2 = Me, R1+R2 = (CH2)5]. Various procedures for synthesizing the phosphinite function were explored in order to improve the yield of the reaction. Results were best when Ph2PNET2 was used in the presence of tetrazol as catalyst. The prepared ligands, which have different electron-donating or electron-withdrawing aryl groups were added to iridium complexes producing catalyst precursors active in the asym. hydrogenation of acetophenone N-benzyl- and N-phenylimines (17, 19). Cationic iridium complexes were more active than the neutral analogs. The use of additives was, in general, detrimental to both the conversion and the enantioselectivity. In the hydrogenation of 17, results were best with ligand 11a (76% ee), but in the hydrogenation of 19 (70% ee) they were best with ligand 10b.
- ST phosphinite phosphite mannitol chiral nonracemic prepn asym hydrogenation catalyst; diarylphosphinite chiral nonracemic bidentate prepn phosphinamidite esterification mannitol; phosphite chiral nonracemic phosphinite prepn mannitol esterification iridium complexation; imine asym hydrogenation catalyst iridium diarylphosphinite phosphite mannitol deriv
- IT Asymmetric synthesis and induction
(asym. hydrogenation; preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)
- IT Phosphorus acids
RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);
USES (Uses)
(esters, phosphinites; preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)
- IT Imines
RL: RCT (Reactant); RACT (Reactant or reagent)
(ketimines; preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)
- IT Phosphites
RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);
USES (Uses)
(preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)
- IT Carbohydrates, reactions
RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)

IT Hydrogenation catalysts
(stereoselective, asym.; preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)

IT 12112-67-3 35138-23-9 666825-73-6
RL: CAT (Catalyst use); USES (Uses)
(preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)

IT 666826-00-2P 666826-05-7P 666826-06-8P 666826-22-8P 881994-91-8P
RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation); USES (Uses)
(preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)

IT 1749-19-5 13685-91-1 13685-97-7 14428-98-9 106054-14-2
110814-25-0 666825-71-4 666825-96-3
RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)

IT 666826-33-1P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)

IT 17480-69-2P 21232-36-0P 21232-37-1P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of chiral C2-sym. bis-phosphinites and C1-sym. phosphinite-phosphites with carbohydrate-derived D-mannitol backbone as ligands for iridium-catalyzed asym. hydrogenation of imines)

RE.CNT 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE

- (1) Aghmiz, M; J Org Chem 2004, V69, P7502 HCAPLUS
- (2) Anon; Catalytic Asymmetric Synthesis 1993
- (3) Anon; Comprehensive Asymmetric Catalysis 1999, V1
- (4) Barbaro, P; Organometallics 2002, V21, P1430 HCAPLUS
- (5) Bayer, A; Eur J Inorg Chem 2002, P2614 HCAPLUS
- (6) Bianchini, C; Eur J Inorg Chem 2001, P43 HCAPLUS
- (7) Blaser, H; Adv Synth Catal 2003, V345, P103 HCAPLUS
- (8) Blaser, H; Isotopics 2002, V19, P3 HCAPLUS
- (9) Buisman, G; Tetrahedron: Asymmetry 1995, V6, P719 HCAPLUS
- (10) Casalnuovo, A; J Am Chem Soc 1994, V116, P9869 HCAPLUS
- (11) Dieguez, M; Chem Commun 2000, P1607 HCAPLUS
- (12) Dieguez, M; Chem Commun 2001, P1132 HCAPLUS
- (13) Dieguez, M; Chem Eur J 2001, V7, P3086 HCAPLUS
- (14) Dieguez, M; J Org Chem 2002, V67, P3796 HCAPLUS
- (15) Dieguez, M; New J Chem 2002, V26, P826
- (16) Guiu, E; Adv Synth Catal 2003, V345, P169 HCAPLUS
- (17) Guiu, E; Tetrahedron: Asymmetry 2004, V15, P3365 HCAPLUS
- (18) Hamamoto, S; Chem Lett 1986, P1401 HCAPLUS
- (19) Holz, J; Eur J Org Chem 2001, V63, P4621
- (20) Horton, D; Carbohydr Res 1973, V30, P367 HCAPLUS
- (21) Kobayashi, S; Chem Rev 1999, V99, P1069 HCAPLUS
- (22) Li, W; J Org Chem 2000, V65, P3489 HCAPLUS
- (23) Li, W; J Org Chem 2002, V67, P7618 HCAPLUS
- (24) Li, W; Tetrahedron Lett 1999, V40, P6701 HCAPLUS
- (25) Margalef-Catala, R; Tetrahedron: Asymmetry 2000, V11, P1469 HCAPLUS
- (26) Morimoto, T; Synlett 1995, P748 HCAPLUS

- (27) Morimoto, T; Tetrahedron: Asymmetry 1995, V6, P2661 HCAPLUS
(28) Otero, D; Carbohydr Res 1984, V128, P79 HCAPLUS
(29) Palmer, M; Tetrahedron: Asymmetry 1999, V10, P2045 HCAPLUS
(30) Pamies, O; Chem Commun 2000, V12, P2383
(31) Pamies, O; J Org Chem 2001, V66, P8364 HCAPLUS
(32) Pfaltz, A; Chimia 2001, V55, P708 HCAPLUS
(33) RajanBabu, T; J Am Chem Soc 1992, V114, P6265 HCAPLUS
(34) RajanBabu, T; J Am Chem Soc 1994, V116, P4101 HCAPLUS
(35) RajanBabu, T; J Am Chem Soc 2001, V123, P10207 HCAPLUS
(36) Rajanbabu, T; J Org Chem 1997, V62, P6012 HCAPLUS
(37) Reetz, M; Tetrahedron: Asymmetry 1999, V10, P2129 HCAPLUS
(38) Sablong, R; Tetrahedron Lett 1996, V37, P4937 HCAPLUS
(39) Selke, R; J Mol Catal 1986, V37, P213 HCAPLUS
(40) Selke, R; J Mol Catal 1993, V84, P223 HCAPLUS
(41) Spindler, F; Angew Chem Int Ed Engl 1990, V29, P558
(42) Spindler, F; Transition Metals for Organic Synthesis 1998, V2, P69 HCAPLUS
(43) Tanaka, T; Tetrahedron Lett 1986, V27, P199 HCAPLUS
(44) Tani, K; Chem Lett 1995, P955 HCAPLUS
(45) Tararov, V; Tetrahedron: Asymmetry 1999, V10, P4009 HCAPLUS
(46) Togni, A; J Am Chem Soc 1994, V116, P4062 HCAPLUS
(47) van Straten, N; Tetrahedron 1997, V53, P6523 HCAPLUS
(48) van Straten, N; Tetrahedron 1997, V53, P6539 HCAPLUS
(49) Vargas, S; Tetrahedron Lett 2005, V46, P2049 HCAPLUS
(50) von Matt, P; Angew Chem Int Ed Engl 1993, V32, P566
(51) Xiao, D; Angew Chem Int Ed 2001, V40, P3425 HCAPLUS
(52) Yan, Y; J Org Chem 2000, V65, P900 HCAPLUS
(53) Yan, Y; Org Lett 2000, V2, P4137 HCAPLUS
(54) Zhu, G; Tetrahedron: Asymmetry 1998, V9, P2415 HCAPLUS

=> d l10 ibib hitstr abs 2-19

L10 ANSWER 2 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:1053108 HCAPLUS

DOCUMENT NUMBER: 143:460341

TITLE: Observation of a 1,5-silyl-migration on fructose

AUTHOR(S): Furegati, Stefan; White, Andrew J. P.; Miller, Andrew D.

CORPORATE SOURCE: Imperial College Genetic Therapies Centre, Department of Chemistry, Imperial College London, London, SW7 2AZ, UK

SOURCE: Synlett (2005), (15), 2385-2387

CODEN: SYNLES; ISSN: 0936-5214

PUBLISHER: Georg Thieme Verlag

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 143:460341

IT 869203-03-2P


RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(crystal structure of; unexpected base-assisted 1,5-silyl migration in fructose resulting in a sterically more crowded product)

RN 869203-03-2 HCAPLUS

CN β -D-Fructofuranose, 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



AB During synthetic studies involving fructose, an unexpected silyl migration was observed, resulting in a sterically more crowded product. 1,4-Silyl migrations have been observed previously taking place in several different carbohydrate derivs. However, here we report for the first time an apparent base-assisted 1,5-silyl migration in fructose, identified by evidence from X-ray crystallog. and 2D-NMR spectroscopy. This novel migration is related to the Brook rearrangement, and appears to be mediated via an anionic, cyclic transition state involving pentavalent silicon.

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 3 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:795363 HCAPLUS

DOCUMENT NUMBER: 142:6223

TITLE: C2-Symmetric Diphosphinite Ligands Derived from Carbohydrates. The Strong Influence of Remote Stereocenters on Asymmetric Rhodium-Catalyzed Hydrogenation

AUTHOR(S): Aghmiz, Mohamed; Aghmiz, Ali; Diaz, Yolanda; Masdeu-Bulto, Anna; Claver, Carmen; Castillon, Sergio
CORPORATE SOURCE: Departament de Quimica Analitica i Quimica Organica, Facultat de Quimica, Universitat Rovira i Virgili, Tarragona, 43005, Spain

SOURCE: Journal of Organic Chemistry (2004), 69(22), 7502-7510
CODEN: JOCEAH; ISSN: 0022-3263

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 142:6223

IT 303764-33-2P 666825-71-4P 797043-20-0P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of C2-sym. diphosphinite ligands derived from carbohydrates for asym. rhodium-catalyzed hydrogenation)

RN 303764-33-2 HCAPLUS

CN L-Iditol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]- (9CI)
(CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 666825-71-4 HCAPLUS

CN D-Mannitol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).




RN 797043-20-0 HCAPLUS

CN D-Mannitol, 2,5-anhydro-1,6-bis-O-[tris(1-methylethyl)silyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



GI



AB Modular ligands of C2 symmetry I-III [R = OCPH₃, OSiMe₂CMe₃, OTs, H, OSi(CHMe₂)₃] were easily prepared from D-glucosamine, D-glucitol, and tartaric acid, resp. The application of these ligands in the rhodium-catalyzed hydrogenation of Me acetamidoacrylate, Me acetamidocinnamate, and di-Me itaconate shows that both the configuration and the substituents at positions 2 and 5 of the THF backbone have a strong influence on the enantioselectivity of the processes.

REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 4 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:746141 HCAPLUS

DOCUMENT NUMBER: 141:395724

TITLE: Facile conversion of O-silyl protected sugars into their corresponding formates using POCl₃·DMF complex

AUTHOR(S): Andrade, Marta M.; Barros, M. Teresa

CORPORATE SOURCE: Faculdade de Ciencias e Tecnologia, Departamento de Quimica, REQUIMTE/CQFB, Universidade Nova de Lisboa, Caparica, 2829-516, Port.

SOURCE: Tetrahedron (2004), 60(41), 9235-9243

CODEN: TETRAB; ISSN: 0040-4020

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 141:395724

IT 303779-98-8P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(facile conversion of O-silyl protected sugars into their corresponding formates using Vilsmeier-Haack complex, POCl₃·DMF)

RN 303779-98-8 HCAPLUS

CN α-D-Glucopyranoside, 1-O-[(1,1-dimethylethyl)dimethylsilyl]-6-O-[(1,1-dimethylethyl)diphenylsilyl]-β-D-fructofuranosyl 6-O-[(1,1-dimethylethyl)diphenylsilyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



AB The direct O-formylation of two selectively protected sugar derivs. using the Vilsmeier-Haack (V-H) complex POCl₃·DMF was studied. Primary O-TBDMS and O-TBDPS ethers of sucrose, the most common disaccharide, underwent regio- and chemoselective O-formylation with this formylating agent. This conversion was also studied with a monosaccharide analog.

REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 5 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:213308 HCAPLUS

DOCUMENT NUMBER: 140:253716

TITLE: Preparation of chiral monophosphorus compounds and their transition metal complexes as catalysts for stereoselective hydrogenation

INVENTOR(S): Meseguer, Benjamin; Militzer, Hans-Christian;

Castillon, Sergio; Claver, Carmen; Guiu, Ester

PATENT ASSIGNEE(S): Bayer Chemicals A.-G., Germany; Lanxess Deutschland GmbH

SOURCE: Eur. Pat. Appl., 31 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1398319	A1	20040317	EP 2003-19803	20030830
EP 1398319	B1	20051109		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK


DE 10242351	A1	20040318	DE 2002-10242351	20020912
AT 309255	E	20051115	AT 2003-19803	20030830
CN 1496991	A	20040519	CN 2003-160281	20030911
US 2004127430	A1	20040701	US 2003-660150	20030911
PRIORITY APPLN. INFO.:			DE 2002-10242351	A 20020912
OTHER SOURCE(S):		CASREACT 140:253716; MARPAT 140:253716		
IT 666826-33-1P				

RL: CAT (Catalyst use); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
(preparation of chiral monophosphorus compds. and their transition metal complexes as catalysts for stereoselective hydrogenation of enamides)

RN 666826-33-1 HCAPLUS

CN D-Mannitol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]-, 3-[bis(2,4-dimethylphenyl)phosphinite] (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




IT 666825-71-4P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of chiral monophosphorus compds. and their transition metal complexes as catalysts for stereoselective hydrogenation of enamides)


RN 666825-71-4 HCAPLUS

CN D-Mannitol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



GI



AB The preparation of monophosphorus compds. I (X = O, bond; R1, R2 = same or different organosilyl; R3, R4 = same or different alkyl, organoamino, organoalkoxy, C2-4 alkylene, arylene, cyclohexylene, ferrocenylene, etc.; R5 = H, C1-20 alkyl, C4-24 aryl, C5-25 arylalkyl, C1-20 haloalkyl, etc.), useful as cocatalyst for transition metal complex catalyzed stereoselective hydrogenation, is described. Thus, preparation of chiral monophosphorus compound II is described starting from 2,5-anhydro-D-mannitol; rhodium/II complex catalyzed stereoselective hydrogenation of $\text{PhC}(\text{:CH}_2)\text{NHCOMe}$ is also described.

REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 6 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:177965 HCAPLUS

DOCUMENT NUMBER: 140:235900

TITLE: Preparation of chiral diphosphines and their transition metal complexes and their use in asymmetric synthesis

INVENTOR(S): Meseguer, Benjamin; Militzer, Hans-Christian; Castillon, Sergio; Claver, Carmen; Diaz, Yolanda; Aghmiz, Mohamed; Guiu, Esther; Aghmiz, Ali; Masdeu, Anna

PATENT ASSIGNEE(S): Bayer A.-G., Germany

SOURCE: Ger. Offen., 34 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10241256	A1	20040304	DE 2002-10241256	20020906
EP 1400527	A1	20040324	EP 2003-18221	20030811
EP 1400527	B1	20060322		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
AT 321059	E	20060415	AT 2003-18221	20030811
US 2005080047	A1	20050414	US 2003-643552	20030819
JP 2004161741	A2	20040610	JP 2003-208112	20030820
CN 1493576	A	20040505	CN 2003-158087	20030821

PRIORITY APPLN. INFO.: DE 2002-10238115 IA 20020821
DE 2002-10241256 A 20020906

OTHER SOURCE(S): CASREACT 140:235900; MARPAT 140:235900

IT 666826-33-1P

RL: CAT (Catalyst use); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)


(preparation of chiral diphosphines and its transition metal complexes and

their use in asym. synthesis)

RN 666826-33-1 HCAPLUS

CN D-Mannitol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]-,
3-[bis(2,4-dimethylphenyl)phosphinite] (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



IT 303764-33-2P 666825-71-4P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)

(preparation of chiral diphosphines and its transition metal complexes and
their use in asym. synthesis)

RN 303764-33-2 HCAPLUS

CN L-Iditol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]- (9CI)
(CA INDEX NAME)


Absolute stereochemistry. Rotation (-).




RN 666825-71-4 HCAPLUS

CN D-Mannitol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]-
(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



GI



I

AB The present invention concerns the preparation of chiral diphosphines their transition metal complexes, and use of complexes in asym. syntheses. Thus, preparation of 2,3-bis-O-(diphenylphosphino)-1,6-dideoxy-2,5-anhydro-D-mannitol I, prepared from 1,6-dideoxy-2,5-anhydro-D-mannitol, and [Rh(cod)₂]BF₄/I catalyzed enantioselective hydrogenation of CH₂:C(NHAc)(CO₂Me) is described.

L10 ANSWER 7 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:808828 HCAPLUS

DOCUMENT NUMBER: 138:187980

TITLE: Facilely accessible multidrug resistance modulator derived from sucrose

AUTHOR(S): Murakami, Nobutoshi; Tamura, Satoru; Iwata, Etsuko; Aoki, Shunji; Akiyama, Shin-ichi; Kobayashi, Motomasa
CORPORATE SOURCE: Graduate School of Pharmaceutical Sciences, Osaka University, Osaka, 565-0871, Japan

SOURCE: Bioorganic & Medicinal Chemistry Letters (2002), 12(22), 3267-3270

CODEN: BMCLE8; ISSN: 0960-894X

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 138:187980


IT 81086-97-7P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and multidrug resistance modulation on KB human cell lines of isovaleryl sucrose derivs.)

RN 81086-97-7 HCAPLUS

CN α-D-Glucopyranoside, 1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]-
β-D-fructofuranosyl 6-O-[(1,1-dimethylethyl)diphenylsilyl]- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



AB Exploration for new MDR-modulators utilizing atractysucroses as scaffolds disclosed 2,3,4,3',4'-O-pentaisovalerylsucrose (I) as a readily accessible medicinal lead. This lead was prepared from sucrose in 65% total yield for three steps. In addition, I exhibited more potent MDR modulating activity than verapamil, a representative modulator of MDR mediated by P-gp.

REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 8 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:612961 HCAPLUS

DOCUMENT NUMBER: 133:335447

TITLE: Synthesis and Conformational Studies of Peptidomimetics Containing Furanoid Sugar Amino Acids and a Sugar Diacid

AUTHOR(S): Chakraborty, T. K.; Ghosh, S.; Jayaprakash, S.; Sharma, J. A. R. P.; Ravikanth, V.; Diwan, P. V.; Nagaraj, R.; Kunwar, A. C.

CORPORATE SOURCE: Centre for Cellular and Molecular Biology, Indian Institute of Chemical Technology, Hyderabad, 500 007, India

SOURCE: Journal of Organic Chemistry (2000), 65(20), 6441-6457
CODEN: JOCEAH; ISSN: 0022-3263

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 133:335447

IT 303764-33-2P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(synthesis and conformational studies of peptidomimetics containing furanoid sugar amino acids as)

RN 303764-33-2 HCAPLUS

CN L-Iditol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]- (9CI)
(CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Furanoid sugar amino acids (I) were synthesized and used as dipeptide isosteres to induce interesting turn structures in small linear peptides. They belong to a new variety of designed hybrid structures that carry both amino and carboxyl groups on rigid furanose sugar rings. Four such mols., 6-amino-2,5-anhydro-6-deoxy-D-gluconic acid (Gaa) and its mannonic, idonic (Iaa), and 3,4-dideoxyidonic congeners were synthesized. The synthesis followed a novel reaction path in which an intramol. 5-exo SN2 opening of the hexose-derived terminal aziridine ring in (II) by the γ -benzyloxy oxygen with concomitant debenzoylation occurred during pyridinium dichromate oxidation of the primary δ -hydroxyl group to carboxyl function, leading to the formation of furanoid sugar amino acid frameworks in a single step. Incorporation of these furanoid sugar amino acids into Leu-enkephalin replacing its Gly-Gly portion gave analogs [(III); R = tBuOC(O), H; R1 = OH, H]. Detailed structural anal. of these mols. by CD and various NMR techniques in combination with constrained mol. dynamics (MD) simulations revealed that two of these analogs [III; P = tBuOC(O); R1 = OH; 2R,5R or 2S,5R] have folded conformations composed of an unusual nine-membered pseudo β -turn-like structure with a strong intramol. H-bond between LeuNH \rightarrow sugarC3-OH. This, in turn, brings the two aromatic rings of Tyr and Phe in close proximity, a prerequisite for biol. activities of opioid peptides. The analgesic activities of III (R = tBuOC(O), H; R1 = OH; 2R,5R) determined by mouse hot-plate and tail-clip methods were similar to that of Leu-enkephalin Me ester. The syn disposition of the β -hydroxy-carboxyl motif on the sugar rings appears to be the driving force to nucleate the observed turn structures in some of these mols. Repetition of the motif on both sides of a furanose ring resulted in a novel mol. design of sugar diacid, 2,5-anhydro-D-idaric acid (IV). Bidirectional elongation of the diacid moieties of IV with identical peptide strands led to the formation of a C2-sym. reverse-turn mimetic 12 which displayed a very ordered structure consisting of identical intramol. H-bonds at two ends between LeuNH \rightarrow sugar-OH, the same as in III (R = tBuOC(O), H; R1 = OH; 2R,5R or 2S,5R).

REFERENCE COUNT: 77 THERE ARE 77 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 9 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:602686 HCAPLUS

DOCUMENT NUMBER: 133:335399

TITLE: Fast Galloylation of a Sugar Moiety: Preparation of Three Monogalloylsucroses as References for Antioxidant Activity. A Method for the Selective Deprotection of tert-Butyldiphenylsilyl Ethers


AUTHOR(S): Barros, M. T.; Maycock, C. D.; Sineriz, F.; Thomassigny, C.

CORPORATE SOURCE: Instituto de Biologia Experimental e Tecnologica, Universidade Nova de Lisboa, Oeiras, P-2780-156, Port.

SOURCE: Tetrahedron (2000), 56(35), 6511-6516
CODEN: TETRAB; ISSN: 0040-4020

PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 133:335399
 IT 303779-98-8
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (acetylation; fast galloylation method for preparation of monogalloyl
 sucroses and method for selective deprotection of tert-
 butyldiphenylsilyl ethers)
 RN 303779-98-8 HCAPLUS
 CN α -D-Glucopyranoside, 1-O-[(1,1-dimethylethyl)dimethylsilyl]-6-O-
 [(1,1-dimethylethyl)diphenylsilyl]- β -D-fructofuranosyl
 6-O-[(1,1-dimethylethyl)diphenylsilyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



AB Three protected new gallotannins, namely the 6'-O-(tri-O-methylgalloyl)-2,3,4,6,1',3',4'-hepta-O-acetylsucrose, the 6'-O-(tri-O-methylgalloyl)-2,3,4,6,1',3',4'-hepta-O-benzoylsucrose and the 6,6'-di-O-tert-butylidiphenylsilyl-1'-O-(tri-O-methylgalloyl)-2,3,4,3',4'-penta-O-acetylsucrose have been prepared in 4 short sequences from sucrose. Methods for rapid galloylation have been studied in order to avoid simultaneous acyl transfer reactions. A method for the deprotection of a tert-butylidiphenylsilyl ether using Br in MeOH has been developed which avoids the intramol. migration of a benzoate group.

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 10 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1993:255218 HCAPLUS

DOCUMENT NUMBER: 118:255218

TITLE: Oligosaccharide microscale analysis by circular dichroic spectroscopy: reference spectra for chromophoric D-fructofuranoside derivatives

AUTHOR(S): Ikemoto, Norihiro; Lo, Lee Chiang; Kim, Oak Kyung; Berova, Nikolina; Nakanishi, Koji

CORPORATE SOURCE: Dep. Chem., Columbia Univ., New York, NY, 10027, USA
 SOURCE: Carbohydrate Research (1993), 239, 11-33

CODEN: CRBRAT; ISSN: 0008-6215

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 147694-16-4P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and acetylation of, with bromobenzoyl chloride)

RN 147694-16-4 HCAPLUS

CN α -D-Fructofuranoside, methyl 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-, 4-[3-(4-methoxyphenyl)-2-propenoate], (E)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.



IT 147694-17-5P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and acylation of, with bromobenzoyl chloride)

RN 147694-17-5 HCAPLUS

CN β -D-Fructofuranoside, methyl 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-, 3-[3-(4-methoxyphenyl)-2-propenoate], (E)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.




IT 147672-48-8P 147672-49-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and acylation of, with bromobenzoyl or methoxycinnamoyl chlorides)

RN 147672-48-8 HCAPLUS


CN α -D-Fructofuranoside, methyl 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 147672-49-9 HCAPLUS
 CN β-D-Fructofuranoside, methyl 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 147672-52-4P 147672-53-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation and acylation of, with methoxycinnamoyl chloride)
 RN 147672-52-4 HCAPLUS
 CN α-D-Fructofuranoside, methyl 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-, 4-(4-bromobenzoate) (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 147672-53-5 HCAPLUS
 CN β-D-Fructofuranoside, methyl 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-, 3-(4-bromobenzoate) (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 147672-57-9P 148556-74-5P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 147672-57-9 HCAPLUS
 CN β -D-Fructofuranoside, methyl 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-, 4-(4-bromobenzoate) (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 148556-74-5 HCAPLUS
 CN α -D-Fructofuranoside, methyl 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-, 3-(4-bromobenzoate) (9CI) (CA INDEX NAME)

Absolute stereochemistry.



AB The microscale anal. method, that is being developed in the authors' group for the structure determination of oligosaccharides, yields monosaccharide derivs. bearing two types of chromophores suitable for exciton-coupling, namely, 4-bromobenzoate (λ_{\max} 245 nm) and 4-methoxycinnamate (λ_{\max}

311 nm). Comparison of the circular dichroic (CD) curves of these subunits to those in the reference library allows for the determination of the sugar identities, linkage positions, and the absolute configurations. The 32 possible derivs. of Me α - and β -D-fructofuranosides bearing four chromophores were prepared and their CD spectra recorded. These data serve to extend the CD library, which already encompasses pyranoside derivs. with the gluco-, galacto-, and manno-configurations, and extend the utility of this methodol. to the anal. of fructose-containing oligosaccharides.

L10 ANSWER 11 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1991:62547 HCAPLUS

DOCUMENT NUMBER: 114:62547

TITLE: Sugar chemistry. VII. Periodate oxidation of sucrose derivatives

AUTHOR(S): Badel, Agnes; Descotes, Gerard; Mentech, Julio

CORPORATE SOURCE: Lab. Chim. Org. II, Univ. Lyon I, Villeurbanne, F-69622, Fr.

SOURCE: Carbohydrate Research (1990), 205, 323-31

CODEN: CRBRAT; ISSN: 0008-6215

DOCUMENT TYPE: Journal

LANGUAGE: French

OTHER SOURCE(S): CASREACT 114:62547


IT 63734-13-4

RL: RCT (Reactant); RACT (Reactant or reagent)
(periodate oxidation of)


RN 63734-13-4 HCAPLUS

CN α -D-Glucopyranoside, 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-
 β -D-fructofuranosyl 6-O-[(1,1-dimethylethyl)dimethylsilyl]- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



GI



AB The periodate oxidation of sucrose derivs. I (R, R2, R3 = OH, Cl, OPh3, OSiMe2CMe3; R1=H, Ac) is generally selective for the D-glucopyranoside group. A cleavage at the C(2)-C(3) or C(3)-C(4) positions was observed for I (R, R2, R3 = OPh3, OSiMe2CMe3) resp. The periodate oxidation was more complete for all other derivs. with cleavage at both C(2)-C(3) and C(3)-C(4).

L10 ANSWER 12 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1991:7081 HCAPLUS

DOCUMENT NUMBER: 114:7081

TITLE: Preparation of sucrose derivatives as bacteriostatics

INVENTOR(S): Badel, Agnes; Descotes, Gerard; Mentech, Julio; Thiriet, Bernard

PATENT ASSIGNEE(S): Beghin-Say S. A., Fr.

SOURCE: Eur. Pat. Appl., 12 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent


LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 349431	A1	19900103	EP 1989-401860	19890629
EP 349431	B1	19920415		
R: BE, CH, DE, ES, FR, GB, IT, LI, NL				
FR 2633626	A1	19900105	FR 1988-8723	19880629
FR 2633626	B1	19920228		
ES 2036818	T3	19930601	ES 1989-401860	19890629
PRIORITY APPLN. INFO.:			FR 1988-8723	A 19880629
OTHER SOURCE(S):	MARPAT 114:7081			
IT 63734-13-4				
RL: RCT (Reactant); RACT (Reactant or reagent) (periodic oxidation of)				
RN 63734-13-4	HCAPLUS			
CN	α -D-Glucopyranoside, 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]- β -D-fructofuranosyl 6-O-[(1,1-dimethylethyl)dimethylsilyl]- (9CI) (CA INDEX NAME)			

Absolute stereochemistry.



GI



AB The title compds. [I; R1, R2, R3 = OH, halo, acyloxy, hydrocarbylsilyloxy; R4, R5 = CHO, CH(OH)CHO; or R4R5 = CH(OH)CH(OH)CH(OH) (glucose configuration); R6, R7 = CHO, or R6R7 = CH(OH)CH(OH) (fructose configuration); however, when R4R5 = CH(OH)CH(OH)CH(OH) (glucose configuration), R6R7 may not be CH(OH)CH(H) (fructose configuration); also, R1, R2, and R3 may not simultaneously be OH] were prepared 1',6,6-O-Tris(tert-butyldimethylsilyl)sacharose was oxidized with Na metaperiodate in H₂O-CHCl₃ at 10° for 12 h to give 1 [R1 = R2 = R3 = OSiMe₂CMe₃, R4 = CHO, R5 = CH(OH)CH(OH) (fructose configuration)]. Saccharose oxide 6,6'-dipalmitate (preparation given) had min. inhibitory concentration of 0.01 mg/mL against Staphylococcus ATCC 6538P.

L10 ANSWER 13 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1989:24205 HCAPLUS

DOCUMENT NUMBER: 110:24205

TITLE: A novel stereospecific synthesis of 5-amino-1-β-D-fructofuranosylimidazole-4-carboxamide

AUTHOR(S): Grouiller, Annie; Mackenzie, Grahame; Najib, Boubker; Shaw, Gordon; Ewing, David

CORPORATE SOURCE: Inst. Natl. Sci. Appl. Lyon, Villeurbanne, 69621, Fr.
SOURCE: Journal of the Chemical Society, Chemical Communications (1988), (10), 671-2
CODEN: JCCCCAT; ISSN: 0022-4936

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 110:24205

IT 117901-65-2P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and deprotection of)



RN 117901-65-2 HCAPLUS

CN 1H-Imidazole-4-carboxamide, 5-amino-1-[1,4,6-tris-O-[(1,1-dimethylethyl)dimethylsilyl]-β-D-fructofuranosyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



GI



AB A β-D-fructofuranose fused oxazolidine-2-thione was isolated as the silyl derivative I, which when desulfurized and treated with α-amino-α-cyanoacetamide gave the silylated 1-β-D-fructofuranosyl aminoimidazole II (R = SiMe₂CMe₃) which when deblocked with methanolic hydrogen chloride produced 5-amino-β-D-fructofuranosylimidazole-4-carboxamide (II; R = H).

L10 ANSWER 14 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1984:531003 HCAPLUS

DOCUMENT NUMBER: 101:131003

TITLE: Sucrose derivatives and the selective benzylation of the secondary hydroxyl groups of 6,1',6'-tri-O-tritylsucrose

AUTHOR(S): Holzapfel, Cedric W.; Koekemoer, Johannes M.; Marais, Charles F.

CORPORATE SOURCE: Chem. Dep., Rand Afr. Univ., Johannesburg, 2000, S. Afr.

SOURCE: South African Journal of Chemistry (1984), 37(2), 57-61
 CODEN: SAJCDG; ISSN: 0379-4350

DOCUMENT TYPE: Journal


LANGUAGE: English

IT 63734-13-4P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation and acetylation of)

RN 63734-13-4 HCAPLUS

CN α -D-Glucopyranoside, 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-
 β -D-fructofuranosyl 6-O-[(1,1-dimethylethyl)dimethylsilyl]- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.



AB The preparation and 500 MHz ^1H -NMR spectra of a number of sucrose derivs. are described. The assignment of the individual proton resonances in these compds. contributed to the identification of the mono- and dibenzoates obtained by benzylation of 6,1',6'-tri-O-tritylsucrose following regioselective activation of the secondary OH groups by reaction with dibutyltin oxide or bis(tributyltin) oxide.

L10 ANSWER 15 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1982:123146 HCAPLUS

DOCUMENT NUMBER: 96:123146

TITLE: Sucrochemistry. Part XXXI. Synthesis and reactions of tert-butyldiphenylsilyl ethers of sucrose

AUTHOR(S): Karl, Horst; Lee, Cheang Kuan; Khan, Riaz

CORPORATE SOURCE: Group Res. Dev., Tate and Lyle Ltd., Reading, RG6 2BX, UK

SOURCE: Carbohydrate Research (1982), 101(1), 31-8
 CODEN: CRBRAT; ISSN: 0008-6215

DOCUMENT TYPE: Journal


LANGUAGE: English

IT 81086-97-7P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and acylation)

RN 81086-97-7 HCAPLUS

CN α -D-Glucopyranoside, 1,6-bis-O-[(1,1-dimethylethyl)diphenylsilyl]-
 β -D-fructofuranosyl 6-O-[(1,1-dimethylethyl)diphenylsilyl]- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.



AB The reaction of sucrose with 1.1 mol equivalent of tert-butyldiphenylsilyl (t-BDPS) chloride in pyridine in the presence of 4-dimethylaminopyridine gave the crystalline 6'-t-BDPS ether (I) in 49% yield, without recourse to column chromatog. I was transformed into the 4,6,1'-trichloride by using SO₂Cl₂. When the silylation of sucrose was performed with 3 mol equivalent of the reagent, chromatog. gave the crystalline 6,6'-di-t-BDPS ether and the 6,1',6'-tri-t-BDPS ether (II) in yields of 78 and 18.7%, resp. II was obtained as the major product on treatment of sucrose with 4.6 mol equivalent of the silylating reagent. Removal of the silyl protecting-group in 6,6'-di-O-tert-butyldiphenylsilylsucrose hexabenzoate, using Bu₄NF, proceeded smoothly, but with 4→6 migration of the benzoate.

L10 ANSWER 16 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1980:110980 HCAPLUS

DOCUMENT NUMBER: 92:110980

TITLE: The complexing properties of a chiral 18-crown-6 derivative incorporating a 2,5-anhydro-D-mannitol residue. A constitutional and stereochemical means of enhancing complexation

AUTHOR(S): Haslegrave, J. Anthony; Stoddart, J. Fraser; Thompson, David J.

CORPORATE SOURCE: Dep. Chem., Univ. Sheffield, Sheffield, S3 7HF, UK

SOURCE: Tetrahedron Letters (1979), (24), 2279-82

CODEN: TELEAY; ISSN: 0040-4039

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 72536-29-9P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and methylation of)


RN 72536-29-9 HCAPLUS

CN D-Mannitol, 2,5-anhydro-1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



GI



AB The anhydromannitol I (R = Me), prepared by standard procedures (yields 62-96%) from I (R = H), condensed with tetraethylene glycol bis(toluenesulfonate) to give 19% 18-crown-6 derivative II. II formed extremely strong 1:1 complexes with alkali metal cations, NH_4 , and alkylammonium cations. Constitutional and stereochem. factors involved in the complexation, free energies of complexation, and the influence of the cation on the complexation are discussed.

L10 ANSWER 17 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1977:536202 HCAPLUS

DOCUMENT NUMBER: 87:136202

TITLE: tert-Butyldimethylsilyl ethers of sucrose

AUTHOR(S): Franke, Fritz; Guthrie, R. D.

CORPORATE SOURCE: Sch. Sci., Griffith Univ., Nathan, Australia

SOURCE: Australian Journal of Chemistry (1977), 30(3), 639-47

CODEN: AJCHAS; ISSN: 0004-9425

DOCUMENT TYPE: Journal

LANGUAGE: English


IT 63734-13-4

RL: RCT (Reactant); RACT (Reactant or reagent)
(desilylation and methylation of)

RN 63734-13-4 HCAPLUS

CN α -D-Glucopyranoside, 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-
 β -D-fructofuranosyl 6-O-[(1,1-dimethylethyl)dimethylsilyl]- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




IT 63734-13-4P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 63734-13-4 HCAPLUS

CN α-D-Glucopyranoside, 1,6-bis-O-[(1,1-dimethylethyl)dimethylsilyl]-
β-D-fructofuranosyl 6-O-[(1,1-dimethylethyl)dimethylsilyl]- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



AB The tert-butyldimethylsilyl group was used as a blocking group in carbohydrate chemical; its selectivity towards primary hydroxyl groups, in the absence of imidazole, was shown by preparation of derivs. of Me α-D-glucopyranoside and sucrose. Me α-D-glucopyranoside was converted into Me 6-O-tert-butyldimethylsilyl-α-D-glucopyranoside and sucrose to 6,1',6'-tri-O-tert-butyldimethylsilylsucrose. In the presence of excess sucrose, a mixture of 6'-O-tert-butyldimethylsilyl-, 6,6'-O-tert-butyldimethylsilyl- and 6,1',6'-tri-O-tert-butyldimethylsilyl-sucroses was formed.

L10 ANSWER 18 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1963:415945 HCAPLUS
DOCUMENT NUMBER: 59:15945
ORIGINAL REFERENCE NO.: 59:2928h,2929a
TITLE: Sucrose derivatives. II. Some silyl and cyanoethyl
ethers and a heptaacetal
AUTHOR(S): Barker, S. A.; Brimacombe, J. S.; Harnden, M. R.;
Jarvis, J. A.
CORPORATE SOURCE: Univ., Birmingham, UK
SOURCE: Chem. Soc. (1963), (June), 3403-6
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable
IT 18919-51-2, Glucopyranoside, 1,6-bis-O-(tricyclohexylsilyl)- β -
D-fructofuranosyl 6-O-(tricyclohexylsilyl)-, α -D-
894412-26-1, Sucrose, 1',6,6'-tris-O-(tricyclohexylsilyl)-
(preparation of)
RN 18919-51-2 HCAPLUS
CN Glucopyranoside, 1,6-bis-O-(tricyclohexylsilyl)- β -D-fructofuranosyl
6-O-(tricyclohexylsilyl)-, α -D- (8CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 894412-26-1 HCAPLUS
CN Sucrose, 1',6,6'-tris-O-(tricyclohexylsilyl)- (7CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 2-A



AB cf. CA 58, 2496b. Selective substitution of sucrose has been achieved by using chlorotricyclohexylsilane. 1,10-Divinylxydecane with sucrose yielded mainly a heptaacetal. Tri-O-vinylsucrose has been produced by transvinylation of sucrose in tetramethylene sulfone. Octa-O(2-cyanoethyl)sucrose was isolated from the mixture produced by repeated reaction of sucrose with acrylonitrile.

L10 ANSWER 19 OF 19 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1963:415944 HCAPLUS

DOCUMENT NUMBER: 59:15944

ORIGINAL REFERENCE NO.: 59:2928f-h

TITLE: Polynucleotides. I. Synthesis of uridylyl-(3' →

5')-uridine and uridylyl-(3' → 5')-6-azauridine

AUTHOR(S): Hall, Ross H.; Thedford, Roosevelt

CORPORATE SOURCE: Roswell Park Mem. Inst., Buffalo, NY

SOURCE: Journal of Organic Chemistry (1963), 28, 1506-9

CODEN: JOCEAH; ISSN: 0022-3263

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

OTHER SOURCE(S): CASREACT 59:15944

IT 18919-51-2, Glucopyranoside, 1,6-bis-O-(tricyclohexylsilyl)-β-

D-fructofuranosyl 6-O-(tricyclohexylsilyl)-, α-D-


(preparation of)

RN 18919-51-2 HCAPLUS

CN Glucopyranoside, 1,6-bis-O-(tricyclohexylsilyl)-β-D-fructofuranosyl

6-O-(tricyclohexylsilyl)-, α -D- (8CI) (CA INDEX NAME)

Absolute stereochemistry.



GI For diagram(s), see printed CA Issue.

AB 2',5'-Di-O-trityluridine serves as a convenient starting point for the synthesis of phosphate dinucleosides containing uridine. This compound was readily phosphorylated with cyanoethyl phosphate and after removal of the cyanoethyl group the resultant blocked nucleotide (I) was used to phosphorylate 2',3'-isopropylideneuridine and 2,3'-isopropylidene-6-azauridine. After removal of blocking groups, the title compds. (II and III) were isolated in good yield from ionexchange columns.

=> fil home

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

102.48

272.21

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-14.25

-14.25

FILE 'HOME' ENTERED AT 14:51:06 ON 22 SEP 2006